

Anti-Stress Effects of *d*-Limonene and Its Metabolite Perillyl Alcohol

Patrizia A. d'Alessio,¹ Jean-François Bisson,² and Marie C Béné³

Abstract

Stress is closely linked by its biological mechanisms to inflammation and by its consequences to accelerated aging. Stress triggers a hormonal response along the hypothalamus–pituitary–adrenal (HPA) axis, which can disrupt the ortho/parasympathetic balance essential for a harmonious life. Proper nutrition, adequate physical activity, and limiting the harmful influence of stress play important roles in avoiding the development of disease and promoting healthy aging. *d*-Limonene, a monoterpene shown to reduce inflammatory parameters in several pre-clinical and clinical models, could also produce an anti-stress action by altering ortho/parasympathetic parameters as well as central neurotransmitter functions. Here we report on a rat model, where a functional observational battery (FOB) was performed by submitting animals to non-pathological stress. *d*-Limonene or its metabolite perillyl alcohol (POH) were administered *per os* at a dose of 10 mg/kg. FOB tests were performed 1 hr before gavage and then at 60, 120, and 180 min. These tests confirmed the stressed status of control rats fed vehicle. Conversely, a series of parameters were significantly less disturbed in treated rats, who retained a better activity and displayed less signs of stress. These effects were more pronounced and sustained after ingestion of *d*-limonene than POH, suggesting the role of endogeneous metabolization of the terpene. These studies show that *d*-limonene exerts, through its metabolite POH, a significant anti-stress action measurable by behavioral and physiologic parameters under the influence of the nervous system. In addition to its anti-inflammatory effects, a beneficial role as an anti-stress substance could thus be claimed for *d*-limonene used as a dietary supplement.

Introduction

ONE OF THE KEY FACTORS responsible for stress syndromes seems to be an imbalance of the sympathetic and parasympathetic nervous systems. Physiologically, the former acts as an accelerator whereas the latter functions as a brake, yet only the sympathetic nervous system is accessible to conscious alterations. For instance, it is possible to increase heart rate through exercise, but it is almost impossible to reduce it voluntarily, and it is therefore necessary to use drugs to restore the ortho/parasympathetic balance to prevent early cardiovascular fatalities.^{1,2} Stress can also cause a hormonal imbalance in the brain, especially in the hypothalamus–pituitary–adrenal (HPA) axis. Stress (whether physical or psychological) induces neurons in the hypothalamus to release corticotrophin-releasing hormone (CRH), which is then transported to the pituitary gland to initiate the secretion of adrenocorticotrophic hormone (ACTH). The adrenal cortex stimulated by ACTH increases the levels of cortisol, known as “the stress hormone.”³ Cortisol indeed has a beneficial function in the body, but

excessive levels brought by chronic stress can cause a dysregulation of inflammatory parameters. HPA and cortisol hyper-activation have thus been reported to be associated with a variety of health problems, such as chronic inflammatory diseases, hypertension, or depression.^{4–6}

During the course of inflammation, the pro-inflammatory cytokines that are produced by peripheral innate immune cells have an impact on brain functions via neural and humoral pathways. Enhancement of interleukin-6 (IL-6) production induces signs of sickness, including reduction of social exploration, immobility, and body weight loss.⁷ Aging, moreover, exacerbates depressive-like behaviors in response to activation of the peripheral innate immune system.⁸ In this context, to counteract these dysbalances and restore an efficient ortho/parasympathetic function seems interesting. Nonetheless, it is also known that neurotransmitters of the central nervous system regulate acute and chronic stress, and major mood-modulating drugs used to lower anxiety and depression act at this central level. For instance, they inhibit serotonin recapture at the level of brain synapses. In response to stress, the hypothalamus is able to produce, among others, consistent

¹Biopark Cancer Campus, University Paris Sud-11, Villejuif, France.

²ETAP Research Centre, Vandœuvre-lès-Nancy, France.

³Hematology Laboratory, CHU Nantes, Nantes, France.

amounts of dopamine,⁹ noradrenaline, and serotonin, which influence the modulation of motor and psychic outcomes.

Until now, it has been clear that neurotransmitter status in the brain as well as certain brain functions/actions (such as appetite, sleep, memory/learning, and emotion) are modified by different nutrients, food components, nutrition conditions,¹⁰ as well as increased stress.¹¹ In turn, it can be assumed that neurotransmitters participate in inducing the stress defense. Compounds derived from natural substances, mostly edible plants, have been known for centuries as "unusual food," in that they are not expected to be consumed for energy purposes, but rather to prevent or treat infections, inflammation, and mood disorders.¹² These compounds have recently acquired a new status of "alicaments," interesting pharmacological candidates for the development of novel drugs preventing, maintaining, and/or curing many body disabilities, the latter being potentially induced or precipitated by stressful events.^{13–15} When added to diet, these natural compounds can act on the autonomic and/or central nervous systems and help to maintain or restore their function. Plants such as valerian, skullcap, and hops have been recognized to activate vagal tone and bring the sympathetic/parasympathetic systems to a balanced alternation. A compound extracted from orange peel, *d*-limonene, could have an even greater potential, because it has been shown that its metabolite perillyl alcohol ([POH] or perillidic acid) induces a peak of dopamine at 4 hr in the hypothalamus after administration of *d*-limonene.¹¹

Materials and Methods

Drugs

d-Limonene (C₁₀H₁₆) of 97% purity and POH ([*S*]-4-isopropenyl-1-cyclohexenylmethanol; [*S*]-*p*-Mentha-1,8-dien-7-ol) of 98% purity were purchased from Sigma-Aldrich (Saint-Quentin Fallavier, France) and prepared each day as a fresh solution by dissolving in corn oil (Olvea, Saint Léonard, France) as vehicle. Stock solutions were 4 mg/mL.

Animal model

Eighteen Wistar HsdBr/Han female rats (175–200 grams) were obtained from EOPS (Harlan Breeding Centre, Gannat, France). At reception, the rats were labeled and maintained in the standardized conditions of an animal house with

22 ± 2°C temperature and 50 ± 10% hygrometry. Food and water were provided *ad libitum*. The animal house had an inverted light/dark cycle of 12 hr with lights on from 21:00 to 09:00. Rats were allowed a 1-week adaptation period to the laboratory conditions. For the tests, rats were fed with vehicle (corn oil), or 10 mg/kg (2.5 mL/kg) of *d*-limonene or POH in vehicle, thus composing three groups of six rats.

Functional observational battery

To evaluate the potential properties of *d*-limonene on stress, a series of tests was proposed to rats in standardized conditions. The battery of functional tests observed was dubbed the functional observational battery (FOB),^{16,17} which was used previously to assess functional deficits in rats exposed to chemicals or to quantify neurotoxic effects.

All animals were observed carefully by a trained technician who was blinded with respect to the animals' treatment; the same observer was used to evaluate the animals throughout the experiment. All animals were observed 60 min prior to and 60, 120, and 180 min after oral administration of test substances. They were removed from the home cage and placed in a standard arena for observation in a specific red-lit room dedicated to the experiment. Efforts were made to ensure minimal variations in sound level, temperature, humidity, lighting, odors, time of day, and environmental distractions. Rats of the different groups were all handled in the same way and under the same conditions and were tested during the first hours of their daily active phase.

Around 40 parameters were measured in about 7 min for each passage of each rat in the experimental devices. Spontaneous locomotor activity and emotional state (anxiety) were measured by observing de-ambulation in an open arm or an open field. Balance was assessed using devices in which the rat was placed head down. Vision and motor coordination were tested through grip and suspension as the rat was approached and then placed on an anti-gravity device. Audition was evaluated by producing a rattling sound near the ears. Approach and contact were measured by nearing a stylus for tactile stimulation without or with touching, respectively. Pinching allowed appreciation of pain perception, *i.e.*, proprioception. A flashlight was used for visual testing. Temperature changes were measured with a digital thermometer. The parameters recorded were spontaneous locomotor activity,

TABLE 1. VARIATION OF BEHAVIORAL AND PHYSIOLOGICAL EFFECTS OF STRESS AT GIVEN TIMES AFTER ADMINISTRATION OF *D*-LIMONENE OR PERILLYL ALCOHOL

	60 min			120 min			180 min		
	<i>H</i> _(df=2) (P)	P DL vs. V	P PA vs. V	<i>H</i> _(df=2) (P)	P DL vs. V	P PA vs. V	<i>H</i> _(df=2) (P)	P DL vs. V	P PA vs. V
Temperature	0.88 (NS)	NS	NS	5.67 (NS)	NS	NS	7.62 (<0.05)	<0.01	NS
Vocalization	8.08 (<0.05)	<0.01	<0.1	6.8 (<0.05)	<0.05	NS	11.3 (<0.01)	<0.001	<0.1
Toe pinch	6.59 (<0.05)	<0.05	<0.05	1.42 (NS)	NS	NS	5.21 (NS)	NS	NS
Startles	5.38 (NS)	NS	NS	7.2 (<0.05)	<0.05	NS	7.77 (<0.05)	<0.01	NS
Approach	3.8 (NS)	NS	NS	9.9 (<0.01)	<0.01	<0.05	4.4 (NS)	NS	NS
Irritability	11.9 (<0.01)	<0.01	<0.01	11.9 (<0.05)	<0.05	NS	4.2 (NS)	NS	NS
Interest	9.32 (<0.01)	<0.01	<0.1	6.8 (<0.05)	<0.05	NS	6.6 (<0.05)	<0.05	NS
Inversion	5.36 (NS)	NS	NS	4.01 (NS)	NS	NS	10.58 (<0.01)	<0.01	NS
Cases	0.5 (NS)	NS	NS	4.25 (NS)	NS	NS	6.12 (<0.05)	<0.05	NS

df, degrees of freedom; DL, *d*-Limonene; PA, perillyl alcohol; V, vehicle; NS, not significant.

body position, movement coordination, head-flicking, head-searching, circling, freezing, backward walking, irritability, tremors, vocalization, lacrimation, salivation, piloerection, pupillary reflex, palpebral reflex, position of the tail, pelvic elevation, and muscle tone of the limbs and abdomen. In the inversion and anti-gravity response tests, forelimb/hind limb grip strength was measured. Upon stimulation, the toe and tail pinch reflexes, visual placing response, and startle response (Preyer reflex) were evaluated. Miction, defecation and consistency of feces, body temperature, and heart and breathing rates were also measured in all conditions.

Classification of studied variables

The variables measured can be categorized according to the functions assessed.

- **Behavioral effects:** Spontaneous locomotive activity, locomotive behavior troubles, anxiety, touch response, irritability, aggression, freezing, somnolence, number of defecations, number of mictions, sensor-motor responses (toe pinch and sound response).
- **Neurological effects:** Pupillary reflex, palpebral closure, pelvic elevation, tail position, limb and abdominal tones, reversal test, grip test, tremors, and piloerection.
- **Physiological effects:** Salivation, lacrimation, diarrhea, body temperature, respiratory rhythm.

Statistical analyses

The variables measured were scored on adapted scales for each parameter. Statistical analyses were carried out using the Statview 5 statistical package (SAS, Institute Inc., USA) and MedCalc Software (Mariakerke, Belgium). Non-parametric tests applied were one-way analysis of variance (ANOVA) with Kruskal–Wallis test followed, when significant, by the Mann–Whitney U-test to compare the different study variables. For all comparisons, differences were considered to be significant at the level of $p < 0.05$.

Results

Data collected during the observation or manipulation of the rats before and after gavage were compared between the control group and treated animals at each time point. The evolution of these parameters over time in each group was also considered. Of note, at baseline, the three groups of animals had strictly similar FOB results.

Comparison of the treated groups with the vehicle group

Because only mild stress was applied (manipulation, postural adaptation, acoustic and visual stimulation), no significant variation of neurological parameters was observed. However, significant differences were observed for a series of behavioral and physiological effects.

Table 1 summarizes these significant variations, which varied depending on the time of testing. At 60 min, the most important difference was observed for irritability, which was significantly less after ingestion of either *d*-limonene or POH. Similarly, treated rats produced less vocalization and reacted less to toe pinch but showed more interest for objects than rats fed corn oil only. At 120 min, the positive effect of

d-limonene was retained for irritability, vocalization, and interest for objects. Additional significant effects were less fear reaction to approach with both compounds and less startles upon rattling when fed *d*-limonene. Significantly less vocalization and more interest for objects was still observed for *d*-limonene-fed rats at 180 min who also displayed less startles. These rats also performed significantly better in balance tests and crossed more cases (*i.e.*, were more active)




FIG. 1. Evolution over time and comparison to the baseline value (–60 min) of three parameters demonstrating the stress imposed on animals fed only corn oil (vehicle, black line with diamonds), and the better behavior of animals who had received POH (dark grey line with triangles) or *d*-limonene (light grey line with squares). Data are expressed as average ranks of the Kruskal–Wallis test.

than rats who received either vehicle or POH. At 180 min, although both POH- and *d*-limonene-fed rats showed a decrease in temperature over the entire experiment, this drop became significant only for the latter. Of note, none of these variations remained significant for POH-fed rats at 180 min.

Evolution over time compared to baseline

Performance comparison over time confirmed that significant stress was applied to the animals, as demonstrated by the evolution of parameters in vehicle-fed rats (Fig. 1). The latter displayed significantly decreased spontaneous activity and interest for objects, and consistently more time in freezing (*i.e.*, fear) than treated rats. Little significant evolution was seen over time for rats receiving POH (only an increase in cases crossed), and any effect, which usually paralleled that seen in rats having received *d*-limonene, appeared to be transient.

Conversely, a large number of parameters differed significantly from baseline in rats fed *d*-limonene. At variance with control rats for which these parameters remained stable, *d*-limonene-fed rats showed significantly decreasing reaction to approach, toe pinch, or tail pinch, became less and less startled or irritable, and almost stopped vocalizing. They also significantly dropped their body temperature while retaining or even improving their ability to keep balance. Examples of these evolutions are shown in Fig. 2.

Discussion

The studies reported here, performed by administering *d*-limonene or POH to rats submitted to non-pathologic mild stress conditions, complete a previous report on the same experiment.¹³ As published in 2008, substantial changes in movement parameters as well as motivational aspects revealed the influential effect of *d*-limonene and POH on the emotional aspects of stress.

This complementary study shows further data regarding physiologic and behavioral changes induced by the oral administration of *d*-limonene acting on the ortho-parasympathetic system. The capacity of *d*-limonene to reverse stress consists of a decrease in startles or aggressions. Individual susceptibility toward aggressiveness may thus be a matter of stress, because startles and elevated body temperature are signs of incoming depression.¹⁸ Indeed, in the Ristomed trial^{14,15} (performed on healthy 65–85-year-old subjects with high inflammatory scores who benefited from orange peel extract (OPE) supplementation containing a high amount of *d*-limonene), a significant decrease of peripheral IL-6 levels was observed, associated with a statistically significant effect on depressive mood.^{14,15} The pronounced anti-stress effects of *d*-limonene in an animal model, as well as of the OPE dietary supplement in humans, suggests that this compound could be worthwhile in multimodal anti-stress therapy concepts.¹⁹

Several natural compounds may have an effect on stress and thereby condition the functional status of many




FIG. 2. Improved parameters compared to the baseline value (–60 min) and to animals fed only corn oil (vehicle, black line with diamonds), and in rats fed POH (dark grey line with triangles) or *d*-limonene (light grey line with squares). Data are expressed as average ranks of the Kruskal–Wallis test.

organs.²⁰ Plants have traditionally been used to manage stress. The rationale of using buckwheat flowers as an herbal medicine is related to its content in vitamin B1, magnesium, and phosphorus. Passionflower has relaxing and sedative properties that could cure not only stress but also insomnia.²¹ Finally, hawthorn, a bush that produces white flowers and red fruit, is used for its calming and sedative properties, without causing drowsiness.²²

Here we have presented new evidence of the boosting effect of *d*-limonene and POH, contained in OPE, as anti-stress agents. Interestingly, although they showed the same trend, the effects of POH were less significant than those of *d*-limonene.²³ The latter even appeared delayed or persistent over the 3 hr of observation and manipulation of the rats. This suggests that endogenous metabolization of *d*-limonene was responsible for the release of its active metabolite POH over the length of the experiment. This is consistent with reports from Yokogoshi¹¹ demonstrating the ability of POH to induce the release of dopamine and control stress.

Acknowledgment

This study was funded by a grant from the French Ministry of Research to Pr. Patrizia A. d'Alessio, the "Award for Innovative Research," 2005.

Author Disclosure Statement

No competing financial interests exist.

References

- Tsigos C, Chrousos GP. Hypothalamic-pituitary-adrenal axis, neuroendocrine factors and stress. *J Psychosom Res* 2002;53:865–871.
- Kelly JJ, Mangos G, Williamson PM, Whitworth JA. Cortisol and hypertension. *Clin Exp Pharmacol Physiol* 1998;25:51–56.
- Zimecki M, Artym J. The effect of psychic stress on the immune response. *Postepy Hig Med Dosw*. 2004;58:166–175.
- Esch T, Stefano GB, Fricchione GL, Benson H. The role of stress in neurodegenerative diseases and mental disorders. *Neurol Endocrinol Lett*. 2002;23:199–208.
- Faramawi MF, Gustat J, Wildman RP, Rice J, Johnson E, Sherwin R. Cardiovascular Health Study Investigators. Relation between depressive symptoms and common carotid artery atherosclerosis in American persons 65 years of age. *Am J Cardiol* 2007;99:1610–1613.
- Du YJ, Zhang HY, Li B, Wu X, Lv YB, Jin HL, Cao YX, Sun J, Luo QL, Gong WY, Liu BJ, Wu JF, Shi SX, Dong JC. Sputum interleukin-6, tumor necrosis factor- α and salivary cortisol as new biomarkers of depression in lung cancer patients. *Prog Neuropsychopharmacol Biol Psychiatry* 2013;47C:69–76.
- Dantzer R, Bluthé RM, Michaud B, Poli V. Role of IL-6 in cytokine-induced sickness behavior: A study with IL-6 deficient mice. *Physiol Behav* 2000;70:367–373.
- Godbout JP, Moreau M, Lestage J, Chen J, Sparkman NL, O'Connor J, Castanon N, Kelley KW, Dantzer R, Johnson RW. Aging exacerbates depressive-like behavior in mice in response to activation of the peripheral innate immune system. *Neuropsychopharmacology* 2008;33:2341–2351.
- Lammel S, Lim BK, Malenka RC. Reward and aversion in a heterogeneous midbrain dopamine system. *Neuropharmacology* 2013; 00115-9 e-pub ahead of print.
- Aoyama M, Sakai T, Yokogoshi H, Yamamoto S. Effect of dietary tyrosine supplementation on the resistance to acute stress in rats. *Jpn J Physiol* 1995;45:S288.
- Nestle Nutrition Council, Japan March 2007, *Nutr Rev*. <http://www.nncj.nestle.co.jp/en/scientificpublications/repositoryofnutritionreviews/nutritionandbrain> Last accessed March 30, 2014.
- Masi S, Gustafsson E, Saint Jalme M, Narat V, Todd A, Bomsel MC, Krief S. Unusual feeding behavior in wild great apes, a window to understand origins of self-medication in humans: Role of sociality and physiology on learning process. *Physiol Behav*. 2012;105:337–49.
- Bisson JF, Menut C, d'Alessio P. Anti-inflammatory senescence activates 5203-L molecule to promote healthy aging and prolongation of lifespan. *Rejuvenation Res* 2008;11:399–407.
- d'Alessio P, Ostan R, Valentini L, Bourdel-Marchasson I, Pinto A, Buccolini F, Franceschi C, Bené MC. Gender differences in response to dietary supplementation by orange peel extract in elderly people in the RISTOMED study: Impact on quality of life and inflammation. *PRIME* 2012;2:30–37.
- d'Alessio PA, Ostan R, Bisson JF, Schulzke JD, Ursini MV, Bené MC. Oral administration of *d*-Limonene controls inflammation in rat colitis and displays anti-inflammatory properties as diet supplementation in humans. *Life Sci* 2013;92:1151–1156.
- MacPhail RC. Observational batteries and motor activity. *Zentralbl Bakteriell Mikrobiol Hyg B* 1987;185:21–27.
- Moser VC. Functional assays for neurotoxicity testing. *Toxicol Pathol* 2011;39:36–45.
- Szuba MP, Guze BH, Baxter LR Jr. Electroconvulsive therapy increases circadian amplitude and lowers core body temperature in depressed subjects. *Biol Psychiatry* 1997; 42:1130–1137.
- d'Alessio P, Ostan R, Capri M, Franceschi C. On the way of longevity: How to counteract neurodegenerative disease. In: Wang Z, Inuzuka H, ed. *Senescence and Senescence-Related Disorders*. INTECH Open Access 2013;6:135–149.
- d'Alessio P, Bennaceur A, Ostan R, Franceschi C. New targets for the identification of an anti-inflammatory anti-senescence activity. In: Nagata T, ed. *Senescence*. Available at <http://tinyurl.com/cpf803> Accessed September 30, 2013.
- Da Silva JDT, Gravena RA, Marques RH, Silva VK, Hada, FH, Barbosa de Moraes VM, Malheiros RD. Passionflower supplementation in diets of Japanese quails at rearing and laying periods. *Revista Brasileira de Zootecnia* 2010;39:1530–1537.
- Atzei AD. *Le piante nella tradizione popolare della Sardegna*. Carlo Delfino editore, Sassari, Italy, 2005.
- Fukumoto S, Sawasaki E, Okuyama S, Miyake Y, Yokogoshi H. Flavor components of monoterpenes in citrus essential oils enhance the release of monoamines from rat brain slices. *Nutr Neurosci* 2006;9:73–80.

Address correspondence to:

Patrizia A. d'Alessio
Biopark Cancer Campus
I, mail Pr G. Mathé 94807 Villejuif France
University Paris Sud-11
CHU Paul Brousse
Avenue Paul Vaillant-Couturier
94807 Villejuif, France

E-mail: patrizia.d-alessio@inserm.fr;
endocell@wanadoo.fr

Received: October 14, 2013

Accepted: October 14, 2013

This article has been cited by:

1. Mi Hee Kim, Jin Il Kim, Eun Ha. 2014. Effects of Aroma-Necklace Application on Perceived Stress, Symptoms of Stress and Changes in Autonomic Nervous System among Nursing Students in Clinical Training. *Journal of Korean Biological Nursing Science* **16**, 334-341. [[CrossRef](#)]